

RESEARCH ARTICLE

Knowledge, attitudes, and practice on the prevention of central line-associated bloodstream infections among nurses in oncological care: A cross-sectional study in an area of southern Italy

Maria Rosaria Esposito¹, Assunta Guillari², Italo Francesco Angelillo^{2*}

1 Istituto Nazionale Tumori, “Fondazione G. Pascale”, Naples, Italy, **2** Department of Experimental Medicine, University of Campania “Luigi Vanvitelli”, Naples, Italy

* italof.angelillo@unicampania.it



OPEN ACCESS

Citation: Esposito MR, Guillari A, Angelillo IF (2017) Knowledge, attitudes, and practice on the prevention of central line-associated bloodstream infections among nurses in oncological care: A cross-sectional study in an area of southern Italy. PLoS ONE 12(6): e0180473. <https://doi.org/10.1371/journal.pone.0180473>

Editor: Lamberto Manzoli, Università degli Studi di Ferrara, ITALY

Received: March 4, 2017

Accepted: June 15, 2017

Published: June 30, 2017

Copyright: © 2017 Esposito et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the paper as Supporting Information files.

Funding: The authors received no specific funding for this work.

Competing interests: The authors have declared that no competing interests exist.

Abstract

The objectives of the cross-sectional study were to delineate the knowledge, attitudes, and behavior among nurses regarding the prevention of central line-associated bloodstream infections (CLABSIs) and to identify their predisposing factors. A questionnaire was self-administered from September to November 2011 to nurses in oncology and outpatient chemotherapy units in 16 teaching and non-teaching public and private hospitals in the Campania region (Italy). The questionnaire gathered information on demographic and occupational characteristics; knowledge about evidence-based practices for the prevention of CLABSIs; attitudes towards guidelines, the risk of transmitting infections, and hand-washing when using central venous catheter (CVC); practices about catheter site care; and sources of information. The vast majority of the 335 nurses answered questions correctly about the main recommendations to prevent CLABSIs (use sterile gauze or sterile transparent semi-permeable dressing to cover the catheter site, disinfect the needleless connectors before administer medication or fluid, disinfect with hydrogen peroxide the catheter insertion site, and use routinely anticoagulants solutions). Nurses aged 36 to 50 years were less likely to know these main recommendations to prevent CLABSIs, whereas this knowledge was higher in those who have received information about the prevention of these infections from courses. Nurses with lower education and those who do not know two of the main recommendations on the site’s care to prevent the CLABSIs, were more likely to perceive the risk of transmitting an infection. Higher education, attitude toward the utility allow to dry antiseptic, and the need of washing hands before wearing gloves for access to port infusion were predictors of performing skin antiseptic and aseptic technique for dressing the catheter insertion site. Educational interventions should be implemented to address the gaps regarding knowledge and practice regarding the prevention of CLABSIs and to ensure that nurses use evidence-based prevention interventions.

Introduction

The issue of Health Care-Associated Infections (HAIs) continues to be one of the most important public health problems in many countries throughout the world [1,2], and these infections remain a leading cause of morbidity and mortality among hospitalized patients [3] also with an increase consumption of resources and add to cost [4,5]. Central line-associated bloodstream infections (CLABSIs), the majority related with the use of the central venous catheter (CVC) [6,7], are the most important complications in critical care [8–10] and in cancer settings [11,12]. In particular, in cancer settings, several risk factors have been identified for CLABSIs in patients who undergo chemotherapy such as, for example, a prolonged duration of catheterization, microbial colonization at the insertion site and of the catheter hub, and inadequate care/maintenance of the CVC after insertion [13–15]. Therefore, improving the quality of the health care is a key priority, since reduction of CLABSIs may be achieved through efforts with adherence to appropriate preventive measures in line with evidence-based recommendations [16,17].

Clinical practice guidelines have been published by several international organisms for the prevention of CLABSIs [18–21] that generally include specific actions to be implemented by health care workers (HCWs) who insert and handle a CVC. Among the HCWs, nurses in cancer settings have the most direct and continuous role in performing high-risk CVC procedures and they should be knowledgeable and compliant in the insertion assistance, care, and maintenance of central lines. Therefore, they are well positioned to implement the recommendations and have a unique opportunity to contribute to primary prevention of these infections via evidence-based best practices [22]. Information has been reported in several surveys in different countries regarding the level of knowledge, the attitudes, and the extent to which evidence-based practices are used among the nurses in different setting [23–29]. To improve the appropriate use of CVCs, in-depth knowledge of the issues by nurses is essential and obtaining this information will contribute to develop preventive programs in order to reduce the frequency of CLABSIs in the cancer setting. Therefore, the objective of the present investigation was to delineate the level of knowledge, attitudes, and behavior regarding CVC procedures among a large sample of nurses working in cancer setting in the Italian hospitals. Furthermore, the second objective was to identify the predisposing factors for the knowledge, attitudes, and behavior.

Materials and methods

The study had a cross-sectional design and was carried out in all 16 non-teaching and teaching public and private hospitals whose units utilized CVCs for oncological patients in the geographic area of Avellino (Ariano Irpino, San Giuseppe Moscati), Benevento (Gaetano Rummo), Caserta (Santa Anna and San Sebastiano), Naples (Cardarelli, Evangelico Villa Betanina, Istituto Nazionale Tumori, Cotugno, Monaldi, San Gennaro, Santa Maria della Pietà, Santa Maria delle Grazie, Santobono/Pausillipon, Second University Teaching Hospital, University Federico II Teaching Hospital), and Salerno (San Giovanni di Dio and Ruggi d'Aragona) (Italy).

Recruitment took place from September to November 2011. The target group was all 472 nurses working in oncology and outpatient chemotherapy units, involved in the care of patients with CVC, in the selected hospitals.

The head physician of each hospital was invited to participate with a letter, they were also informed about the voluntary and anonymous participation and the study coordinator, so that they could contact for ask questions, and questionnaires with detailed instructions about the modality of gather the information. Participants were informed about the purposes of the

study, that the data would be handled confidentially, and that the results would be reported at an aggregated group rather than at an individual level. Written informed consent was obtained before the return of the questionnaire. The questionnaire did not contain identifying information and it was self-administered to the sample that agreed to participate. No incentives or financial payments have been provided for the participation in the study.

The questionnaire was developed by the research team, based on the CDC Guidelines for the Prevention of Intravascular Catheter Related Infections [18]. The complete questionnaire in Italian and English are found in [S1 Questionnaire](#) and [S2 Questionnaire](#). The questions were distributed through five parts. The first part recorded data on nurses' demographic and professional characteristics (age, gender, educational level, years of working activity, unit of activity). The second part was designed to assess nurses' knowledge about evidence-based practices for the prevention of CLABSIs, such as frequency of CVC dressing changes, use of sterile gauze or transparent semipermeable dressings to cover the catheter site, antiseptic agent to use in skin for catheter insertion site, use of topical antibiotic ointment at CVC insertion site, flush the lumen after the administration of medication or fluid, replacement of intravenous (IV) administration sets, disinfect IV ports/needleless connectors before accessing, and use routinely anticoagulants solutions. The options of the responses were "yes", "do not know", and "no". The third part was composed on questions measuring awareness towards the utility of guidelines, perception of the risk of transmitting a CLABSI when handling the CVC and necessity of hand hygiene before and after dressing changes of the catheter site, monitor the catheter site visually when changing the dressing or by palpation through an intact dressing on a regular basis, and hands hygiene before wear gloves for access to port infusion. Three questions were scored using 3-point Likert-type scale with options for "agree", "uncertain", and "disagree" and three questions using a 10-point Likert-type scale, ranging from 1 to 10. The fourth part collected information on practices about catheter site care, dressing and aseptic technique for dressing changes of the catheter insertion site, aseptic technique and correct frequency about disinfection of IV access ports and needleless connectors before accessing or manipulation, replacement of IV administration sets, and replacement sets used to administer blood, blood product, and lipid emulsions. The questions were measured using a 5-point Likert-type scale ranging from "never" to "always", and closed-end questions investigated nurses' practice toward the prevention of CLABSIs. The last part queried nurses on their source of information about CLABSIs and about their interest in acquiring new information about prevention of CLABSIs.

The questionnaire was pilot tested with a group of 20 oncology nurses in order to check the clarity and readability of the questionnaire. The final version of the questionnaire was refined and corrected based on feedback from the participants.

Prior to commencement of data collection, ethical approval for the conduction of the research was granted from the Ethics Committee of the Second University of Naples.

Statistical analysis

A univariate analysis was performed with a *t*-test or chi-square test as appropriate in order to assess the significant association between outcomes variables and the explanatory variables. Variables with *p*-value ≤ 0.25 on univariate analysis were considered for possible entry in the multivariate linear and logistic regressions models. Three separate multivariate stepwise logistic and linear regression models were constructed for identification of the independent association with the selected predictors and the following outcomes of interest: knowledge about the main recommendations to prevent CLABSIs (Model 1); perception of the risk of transmitting a CLABSI when handling the CVC (Model 2); and appropriate behavior about skin antiseptic

and aseptic technique for dressing the catheter insertion site (Model 3). In Model 1, nurses were divided in those who knew about the use sterile gauze or sterile transparent semipermeable dressing to cover the catheter site, disinfect the needleless connectors before administer medication or fluid, disinfect with hydrogen peroxide the catheter insertion site, and use routinely anticoagulants solutions versus all others, and in Model 3, nurses were divided in those who had an appropriate behavior about skin antiseptic and aseptic technique for dressing the catheter insertion site versus all others.

The following explanatory variables were included in all models: gender (male = 0; female = 1), age (categories: $\leq 35 = 1$; 36–40 = 2; 41–45 = 3; 46–50 = 4; $> 50 = 5$), educational level in Nursing (Regional Diploma = 0; University Diploma/Degree = 1), number of years in practice (1–5 = 1; 6–10 = 2; 11–15 = 3; 16–20 = 4; $> 20 = 5$), work setting (outpatient chemotherapy unit = 0; oncology ward = 1), workshops and courses as sources of information on the prevention of CLABSIs (no = 0; yes = 1), and need of additional information about the prevention of CLABSIs (no = 0; yes = 1). The following variables were also included: knowledge about the guidelines for the preventions of CLABSIs regarding dressing changes of catheter insertion site (no = 0; yes = 1), skin antiseptic preparations used for dressing replacement (no = 0; yes = 1), routinely use anticoagulant therapy to reduce the risk of CLABSIs (no = 0; yes = 1), and disinfection of IV ports/needleless connectors before access port (no = 0; yes = 1) in Model 1; perceived the risk of transmitting a CLABSI (continuous) in Model 2; appropriate behavior about skin antiseptic and aseptic technique for dressing the catheter insertion site (no = 0; yes = 1), positive attitude toward the utility of allow to dry antiseptic (no = 0; yes = 1), and necessity of hands hygiene before wearing gloves for access to port infusion (no = 0; yes = 1) in Model 3; correct knowledge about the use of the sterile gauze or transparent semipermeable dressing to cover catheter site and change the dressing of the insertion site every 7 days for semipermeable dressing or if the dressing is soiled or loosened (no = 0; yes = 1) in Models 2 and 3.

The stepwise selection procedure with a forward method was used, and a significance level of 0.2 was used as the criterion for variables to enter in the regression models and 0.4 for variables to remain. Results of the logistic regression models are presented as odds ratios (ORs) with 95% confidence intervals (CIs) and *p*-values. Results of the linear regression model are presented as standardized regression coefficients (β). *P*-values of ≤ 0.05 were considered statistically significant, and all analyses were two-sided. All data (S3 Data Set) were analyzed using the statistical software Stata 10.1.

Results

Demographic and professional characteristics

A total of 335 questionnaires were collected back out of 472 that were initially distributed, giving a response rate of 71%. The socio-demographic and professional characteristics of respondents are shown in [Table 1](#). Approximately two-thirds were female (61.3%) with a mean age of 43 years, the vast majority was in oncology wards, and the average number of years working in these wards was 9.9.

Knowledge about evidence-based practice for the prevention of CLABSIs

The survey responses related to knowledge regarding evidence-based practices for the prevention of CLABSIs are reported in [Table 2](#). The vast majority of nurses, with frequencies ranging from 70.7% to 90.1%, answered questions correctly about to flush the lumen with saline after

Table 1. Main socio-demographic characteristics of the responders.

	<i>n</i> *	%
<i>Gender</i>		
Female	203	61.3
Male	128	38.7
<i>Age group, (years)</i>		
≤35	70	21
36–40	72	21.7
41–45	69	20.8
46–50	57	17.2
>50	64	19.3
<i>Number of years in practice</i>		
1–5	131	39.5
6–10	98	29.5
11–15	22	6.6
16–20	49	14.8
>20	32	9.6
<i>Educational level in Nursing</i>		
Regional Diploma	236	70.5
University Diploma/Degree	99	29.5
<i>Professional role</i>		
Ordinary nurse	310	94.5
Head nurse	18	5.5
<i>Work setting</i>		
Oncology ward	284	84.8
Outpatient chemotherapy unit	51	15.2

* Number of responding

+ Mean±Standard deviation (range)

<https://doi.org/10.1371/journal.pone.0180473.t001>

administration of medication or fluid, to use sterile gauze or transparent semipermeable dressing to cover the catheter site, to not use topical antibiotic ointment on insertion site, and to replace IV administration sets every 72 hours. Only 64.4% acknowledged that the routinely

Table 2. Knowledge about evidence-based practices for the prevention of CLABSIs.

	Yes	No	Do not know
	(%)	(%)	(%)
Flush the lumen with normal saline after the administration of medication or fluid	90.1	7.2	2.7
Use sterile gauze or sterile transparent semipermeable dressing to cover the catheter site	85.9	9.3	4.8
Disinfect the needleless connectors before administer medication or fluid	77.5	12.3	10.2
Replace catheter site dressing every 7 days or if the dressing becomes visibly soiled or loosened	76	19.5	4.5
Use topical antibiotic ointment on catheter insertion site	16.5	75.5	8
Replace the IV administration sets every 72 hours	70.7	19.7	9.6
Disinfect with hydrogen peroxide the catheter insertion site	20	70	10
Use routinely anticoagulants solutions	26	64.4	9.6

In bold are indicated the correct answers

<https://doi.org/10.1371/journal.pone.0180473.t002>

use of anticoagulants solutions does not prevent CLABSIs and 70% that hydrogen peroxide is not recommended to disinfect the catheter insertion site.

The results from the multivariate logistic and linear regression models exploring the association between the different variables and the outcomes of interest are shown in Table 3. Nurses in the age groups from 36–40 years (OR = 0.35; 95% CI 0.16–0.76), 41–45 (OR = 0.41; 95% CI 0.19–0.88), and 46–50 (OR = 0.4; 95% CI 0.17–0.94), when the age ≤ 35 years was chosen as reference category, were less likely to know the main recommendations to prevent CLABSIs,

Table 3. Multivariate logistic (1 and 3) and linear (2) regression models results.

Variable	OR	95% CI	p
Model 1. Knowledge about the main recommendations to prevent CLABSIs			
Log likelihood = -155.08, chi-square = 21.04 (5 df), $p = 0.0008$			
Workshops and courses as sources of information about prevention of CLABSIs	2.44	1.29–4.62	0.006
Age			
$\leq 35^*$	1.0	-	-
36–40	0.35	0.16–0.76	0.008
41–45	0.41	0.19–0.89	0.023
46–50	0.4	0.17–0.94	0.035
Need of additional information about the prevention of CLABSIs	1.75	0.73–4.22	0.21
Model 3. Appropriate behavior about skin antiseptic and aseptic technique for dressing the catheter insertion site			
Log likelihood = -164.46, chi-square = 87.08 (7 df), $p < 0.00001$			
Attitude toward the utility allow to dry antiseptic	7.31	3.96–13.49	<0.001
Attitudes toward the necessity of hands hygiene before wearing gloves for access to port infusion	11.85	3.43–40.89	<0.001
Educational level in Nursing	2.11	1.17–3.8	0.012
Age			
$\leq 35^*$	1.0	-	-
41–45	1.41	0.7–2.84	0.33
46–50	0.67	0.33–1.36	0.27
Workshops and courses as sources of information about prevention of CLABSIs	1.43	0.82–2.49	0.21
Knowledge about the use of the sterile gauze or transparent semipermeable dressing to cover catheter site and change the dressing of the insertion site every 7 days or if the dressing is soiled or loosened	1.32	0.71–2.44	0.38
Variable	Coeff.	t	p
Model 2. Perception of the risk of transmitting a CLABSI			
F (2,328) = 7.25, $R^2 = 4.2\%$, adjusted $R^2 = 3.7\%$, $p = 0.0008$			
Knowledge about the use of the sterile gauze or transparent semipermeable dressing to cover catheter site and change the dressing of the insertion site every 7 days or if the dressing is soiled or loosened	-0.85	-2.66	0.008
Educational level in Nursing	-0.85	-2.6	0.01
Constant	7.15	-	-

*Reference category

<https://doi.org/10.1371/journal.pone.0180473.t003>

whereas this knowledge was higher in those who have received information about the prevention of CLABSIs from workshops and courses (OR = 2.44; 95% CI 1.29–4.62) (Model 1).

Attitudes towards the prevention of CLABSIs

Attitudes towards the effectiveness of guidelines and of hand hygiene before and after dressing changes of catheter insertion site, on a scale from 1 to 10, showed a mean score for the whole sample respectively of 9.1 ± 1.5 and 9.7 ± 1 . Respondents indicated their perceived risk of transmitting to a patient a CLABSI when handling the CVC, on a scale from 1 to 10, with a mean score of 6.3 ± 2.8 . Moreover, 81.8% agreed with recommendations about monitoring the catheter site visually or by palpation through an intact dressing on a regular basis, 85.3% disagree that the use of gloves before the infusion port access replaces the need of hands washing, and only 55.2% agreed that is useful to dry the antiseptic on the insertion site before catheter insertion. The multivariate linear regression analysis allowed to identify several variables associated with the dependent variable “perception of the risk of transmitting a CLABSI”. Nurses with a lower level of education and those who do not know two of the main recommendations on the site’s care for the prevention of catheter related infections were more likely to perceive the risk (Model 2 in Table 3).

Practices about the prevention of CLABSIs

Among the different activities related to the care of patients with CVC, the vast majority was involved in catheter site care. Appropriate procedures were self-reported regarding hands hygiene before dressing (84.6%), hands hygiene with antiseptic soap (81.7%), hands washing for more than 1 minute (58%), use of gloves during dressing replacement (93.8%), povidone-iodine for dressing insertion sites (87.8%), and let the antiseptic dry (33.3%). Results of the multivariate logistic regression model carried out with the practices regarding skin antiseptic and aseptic technique for dressing of the insertion catheter site as outcome variable, revealed that nurses with a graduate degree were 2.1 times more likely to have an appropriate behavior than those with a lower education (95% CI 1.17–3.79). Respondents with a better attitude toward the utility allow to dry antiseptic and toward the need of washing hands before wearing gloves for access to port infusion were respectively 7.31 (95% CI 3.96–13.49) and 11.85 (95% CI 3.43–40.88) more likely to perform skin antiseptic and aseptic technique for dressing the catheter insertion site (Model 3 in Table 3). The responses related to the disinfection of IV access ports indicated that 75.3% of the sample self-reported always disinfects the port infusion with povidone iodine (72.8%). A total of 73% and 79% always wash their hands and wear gloves before replacement of IV administrations sets, respectively. Only 60% replace the sets within 24 hours of initiating the infusion with lipid emulsions, 76.9% always disinfect IV access port needleless connectors before accessing or manipulation, but only 27% swabbing with chlorhexidine 2% as antiseptic agent to clean the access hub or connector, and 76.7% always flushed the lumens with normal saline after IV therapy.

Nurses indicated workshops and courses (67.3%) as their main source of information about the prevention of CLABSIs, followed by guidelines (42.7%), and internet (30.7%). The vast majority (83.7%) reported that they would like to learn more.

Discussion

The data of the current study are among the first reporting knowledge, attitudes, and evidence-based practices for the prevention of CLABSIs among a large sample of nurses working in oncology settings in Italy.

This study demonstrated that nurses have an adequate level of knowledge concerning evidence-based recommendations for preventing CLABSIs, since the vast majority were aware about the main recommendations for patients with CVCs, such as the type of dressing (85.9%) and/or frequency changes (76%) of the catheter insertion site. These findings are comparatively higher than the values observed in nurses in the adult ICU with knowledge of 10% [30] and of 43.4% and 26.2% [26], in ICU and outside ICUs with 37.4% and 62% [27], in ICUs and other wards with 63.9% [24], and in neonatal and pediatric ICUs with 16.7% and 71.8% [31]. In contrast, there are wide areas where the knowledge was lower, particularly regarding the agent for the skin antiseptics, since 30% did not know that hydrogen peroxide is not indicated. These findings suggest the need for including the current evidence-based practice guidelines in educational curricula and programs for HCWs to help them in improving their knowledge. This is also supported by the fact that over three-fourths of the nurses reported that they would like to learn more about the prevention of CLABSIs. Moreover, provision of information about the guidelines for the prevention of CLABSIs influences knowledge since nurses were able to answer correctly if they have received information from workshops and courses.

The findings from this survey showed that respondents had an extremely positive attitude towards the utility of guidelines for the prevention of CLABSIs. A similar result has been observed in nurses in surgical wards in Italy about the utility of guidelines and protocols for disinfection procedures for the prevention of HAIs [32]. Positive attitudes were also recorded regarding the necessity of washing hands before and after dressing the insertion site. The multivariable regression analysis showed that nurses with a high level of perceived risk of transmitting a CLABSI were those with a Regional Diploma level of education and who did not know the recommendations for the care of the catheter site.

Although most nurses' claims that guidelines were necessary in order to reduce CLABSIs, hands hygiene is alarmingly low and with the current evidence that the hands represent the main route of transmission of hospital pathogens, hands hygiene is one of the priority measures on health programs and actions destined to prevent infections. Low adherence to hand hygiene (40–72%) on the nurses' practice toward CVC's care has been reported in teaching and non-teaching hospital ICUs in Yemen [33] and only 26% of the sample that washed their hands successfully in Mongolia [34]. Before central venous catheter insertion, only 22.5% of nurses performed hand hygiene in the ICUs in Egypt [30]. Regarding the use of chlorhexidine 2%, only 27% of the sample used it as antiseptic agent to clean the access port or needleless connectors. This is in accordance with similar previous studies [35, 36].

Based on multivariate regression analysis, the knowledge of recommendations on dressing insertion site was significantly associated with the perception of the risk of transmitting a CLABSI when handling the CVC. Moreover, workshops and courses as source of information were significantly associated with a higher level of knowledge about the main recommendations for the prevention of CLABSIs. The importance of continuous training process that involves nurses allows them to acquire new skills for the CVCs management can explain this finding. Furthermore, the analysis of the predictors of the knowledge also showed that those younger were more knowledgeable. A possible explanation is that nurses of younger age are more likely to take advantage from educational level in university compared to those older with different educational path. This is confirmed by the finding that nurses with a graduate degree perform an appropriate behavior about skin antiseptic and aseptic technique for dressing the catheter insertion site.

Cautions needs to be taken while interpreting this data from the questionnaire as some potential limitations associated with the design and measurements need to be addressed. First, as this is a cross-sectional study, any causative relationship between the examined variables and the outcomes of interest was difficult to determine. Second, similar to all data that are based on self-

report questionnaires, the accuracy of the results was heavily dependent on the honesty and understanding of the respondents, there could have been potential for recall bias. Third, as a general limitation to questionnaire, nurses may tend to provide more socially desirable responses which may show appropriate practice but may not reflect reality. To overcome social desirability bias and improve the validity of the data, it has been used an anonymous self-administered questionnaire which may minimise social desirability bias and increase respondents' willingness to participate. Fourth, the study was limited to nurses in oncology units, so the results may not be generalizable to other nurses and HCW populations, although this was not an aim of the survey. In spite of these limitations, the large sample size and the high response rate in the study reduce the likelihood of bias in the sample and this study provides primary yet valuable data.

Conclusions

In conclusion, the results of this study clearly indicated that the educational interventions are very important and should be implemented to address the gaps regarding knowledge and practice and to ensure that nurses use evidence-based prevention interventions.

Supporting information

S1 Questionnaire.

(DOC)

S2 Questionnaire.

(DOC)

S1 Dataset.

(XLS)

Acknowledgments

The authors are grateful to the support of all selected hospitals and units. The authors would like to thank the nurses who participated in this study.

Author Contributions

Conceptualization: Maria Rosaria Esposito, Italo Francesco Angelillo.

Data curation: Maria Rosaria Esposito, Italo Francesco Angelillo.

Formal analysis: Maria Rosaria Esposito, Assunta Guillari, Italo Francesco Angelillo.

Investigation: Maria Rosaria Esposito, Assunta Guillari.

Methodology: Maria Rosaria Esposito, Italo Francesco Angelillo.

Project administration: Maria Rosaria Esposito, Italo Francesco Angelillo.

Resources: Maria Rosaria Esposito, Assunta Guillari, Italo Francesco Angelillo.

Supervision: Italo Francesco Angelillo.

Validation: Maria Rosaria Esposito, Assunta Guillari, Italo Francesco Angelillo.

Visualization: Maria Rosaria Esposito, Assunta Guillari, Italo Francesco Angelillo.

Writing – original draft: Italo Francesco Angelillo.

Writing – review & editing: Italo Francesco Angelillo.

References

1. Vaz LE, Kleinman KP, Kawai AT, Jin R, Kassler WJ, Grant PS, et al. Impact of medicare's hospital-acquired condition policy on infections in safety net and non-safety net hospitals. *Infect Control Hosp Epidemiol*. 2015; 36: 649–55. <https://doi.org/10.1017/ice.2015.38> PMID: 25732568
2. AHRQ Projects to Prevent Healthcare-Associated Infections, Fiscal Year 2011. October 2014. Agency for Healthcare Research and Quality, Rockville, MD. Available from: <http://www.ahrq.gov/sites/default/files/publications/files/haify11.pdf>. Accessed October 1, 2015.
3. Centers for Disease Control and Prevention. 2013 National and State Healthcare-Associated Infections Progress Report. Published January 14, 2015. Available from: www.cdc.gov/hai/progress-report/index.html. Accessed November 7, 2014.
4. Nelson RE, Jones M, Liu CF, Samore MH, Evans ME, Graves N, et al. The impact of healthcare-associated methicillin-resistant *Staphylococcus aureus* infections on post-discharge healthcare costs and utilization. *Infect Control Hosp Epidemiol*. 2015; 36: 534–42. <https://doi.org/10.1017/ice.2015.22> PMID: 25715806
5. Krein SL, Fowler KE, Ratz D, Meddings J, Saint S. Preventing device associated infections in US hospitals: national surveys from 2005 to 2013. *BMJ Qual Saf*. 2015; 24: 385–92. <https://doi.org/10.1136/bmjqs-2014-003870> PMID: 25862757
6. The Joint Commission. Preventing Central Line-Associated Bloodstream Infections: a global challenge, a global perspective. Oak Brook, IL: The Joint Commission Resources, May 2012. Available from: https://www.jointcommission.org/assets/1/18/CLABSI_Monograph.pdf. Accessed March 11, 2014.
7. Zing W, Sandoz L, Inan C, Cartier V, Clergue F, Pittet D, et al. Hospital-wide survey of the use of central venous catheters. *J Hosp Infect*. 2011; 77: 304–8. <https://doi.org/10.1016/j.jhin.2010.11.011> PMID: 21288595
8. Kallen AJ, Patel PR, O'Grady NP. Preventing catheter-related bloodstream infections outside the intensive care unit: expanding prevention to new settings. *Clin Infect Dis*. 2010; 51: 335–41. <https://doi.org/10.1086/653942> PMID: 20572762
9. Furuya EY, Dick A, Perencevich EN, Pogorzelska M, Goldmann D, Stone PW. Central line bundle implementation in US intensive care units and impact on bloodstream infections. *PLoS One*. 2011; 6(1): e15452. <https://doi.org/10.1371/journal.pone.0015452> PMID: 21267440
10. Apostolopoulou E, Raftopoulos V, Filintisis G, Kithreotis P, Stefanidis E, Galanis P, et al. Surveillance of device-associated infection rates and mortality in 3 Greek intensive care units. *Am J Crit Care*. 2013; 22: e12–20. <https://doi.org/10.4037/ajcc2013324> PMID: 23635940
11. Hentrich M, Schalk E, Schmidt-Hieber M, Chaberny I, Mousset S, Buchheidt D, et al. Central venous catheter-related infections in hematology and oncology: 2012 updated guidelines on diagnosis, management and prevention by the Infectious Diseases Working Party of the German Society of Hematology and Medical Oncology. *Ann Oncol*. 2014; 25: 936–47. <https://doi.org/10.1093/annonc/mdt545> PMID: 24399078
12. Wang TY, Lee KD, Chen PT, Chen MC, Chen YY, Huang CE, et al. Incidence and risk factors for central venous access port-related infection in Chinese cancer patients. *J Formos Med Assoc*. 2015; 114: 1055–60. <https://doi.org/10.1016/j.jfma.2015.06.013> PMID: 26253646
13. Yacobovich J, Ben-Ami T, Abdalla T, Tamary H, Goldstein G, Weintraub M, et al. Patient and central venous catheter related risk factors for blood stream infections in children receiving chemotherapy. *Pediatr Blood Cancer*. 2015; 62: 471–6. <https://doi.org/10.1002/pbc.25281> PMID: 25327811
14. Tomlinson D, Mermel LA, Ethier MC, Matlow A, Gillmeister B, Sung L. Defining bloodstream infections related to central venous catheters in patients with cancer: a systematic review. *Clin Infect Dis*. 2011; 53: 697–710. <https://doi.org/10.1093/cid/cir523> PMID: 21890775
15. Raad I, Chافتari AM. Advances in prevention and management of Central Line-Associated Bloodstream Infections in patients with cancer. *Clin Infect Dis*. 2014; 59 Suppl 5: S340–3.
16. Chopra V, Krein SL, Olmsted RN, Safdar N, Saint S, et al. Prevention of Central Line-Associated Bloodstream Infections: brief update review. In: *Making Health Care Safer II: An Updated Critical Analysis of the Evidence for Patient Safety Practices*. Rockville (MD): Agency for Healthcare Research and Quality (US); 2013. Chapter 10. Available from: <https://www.ncbi.nlm.nih.gov/books/NBK133364/>. Accessed May 5, 2017.
17. Moureau NL, Flynn J. Disinfection of Needleless Connector Hubs: Clinical Evidence Systematic Review. *Nurs Res Pract*. 2015; 2015: 796762. <https://doi.org/10.1155/2015/796762> PMID: 26075093
18. O'Grady NP, Alexander M, Burns LA, Dellinger EP, Garland J, Heard SO, et al. Guidelines for the prevention of intravascular catheter-related infections. *Am J Infect Control*. 2011; 39(4 Suppl 1): S1–34.

19. Schiffer CA, Mangu PB, Wade JC, Camp-Sorrell D, Cope DG, El-Rayes BF, et al. Central venous catheter care for the patient with cancer: American Society of Clinical Oncology clinical practice guideline. *J Clin Oncol*. 2013; 31: 1357–70. <https://doi.org/10.1200/JCO.2012.45.5733> PMID: 23460705
20. Loveday HP, Wilson JA, Pratt RJ, Golsorkhi M, Tingle A, Bak A, et al. epic3: national evidence-based guidelines for preventing healthcare-associated infections in NHS hospitals in England. *J Hosp Infect*. 2014; 86: Suppl 1: S1–70.
21. Ling ML, Apisarnthanarak A, Jaggi N, Harrington G, Morikane K, Thu le TA, et al. APSIC guide for prevention of Central Line Associated Bloodstream Infections (CLABSI). *Antimicrob Resist Infect Control*. 2016; 5: 16. <https://doi.org/10.1186/s13756-016-0116-5> PMID: 27152193
22. Flodgren G, Conterno LO, Mayhew A, Omar O, Pereira CR, Shepperd S. Interventions to improve professional adherence to guidelines for prevention of device-related infections. *Cochrane Database Syst Rev*. 2013; (3): Cd006559. <https://doi.org/10.1002/14651858.CD006559.pub2> PMID: 23543545
23. Ullman AJ, Long DA, Rickard CM. Paediatric ICU nurses: preventing central venous device infections. *Br J Nurs*. 2014; 23: S14–5.
24. Bianco A, Coscarelli P, Nobile CG, Pileggi C, Pavia M. The reduction of risk in central line-associated bloodstream infections: knowledge, attitudes, and evidence-based practices in health care workers. *Am J Infect Control*. 2013; 41: 107–12. <https://doi.org/10.1016/j.ajic.2012.02.038> PMID: 22980513
25. Koutzavekiaris I, Vouloumanou EK, Gourni M, Rafailidis PI, Michalopoulos A, Falagas ME. Knowledge and practices regarding prevention of infections associated with central venous catheters: a survey of intensive care unit medical and nursing staff. *Am J Infect Control*. 2011; 39: 542–7. <https://doi.org/10.1016/j.ajic.2010.11.003> PMID: 21496955
26. Labeau SO, Vandijck DM, Rello J, Adam S, Rosa A, Wenisch C, et al. Centers for Disease Control and Prevention guidelines for preventing central venous catheter-related infection: results of a knowledge test among 3405 European intensive care nurses. *Crit Care Med*. 2009; 37: 320–3. <https://doi.org/10.1097/CCM.0b013e3181926489> PMID: 19050628
27. Dedunska K, Dyk D. Prevention of central venous catheter-associated bloodstream infections: A questionnaire evaluating the knowledge of the selected 11 evidence-based guidelines by Polish nurses. *Am J Infect Control*. 2015; 43: 1368–71. <https://doi.org/10.1016/j.ajic.2015.07.022> PMID: 26307045
28. Shapey IM, Foster MA, Whitehouse T, Jumaa P, Bion JF. Central venous catheter-related bloodstream infections: improving post-insertion catheter care. *J Hosp Infect*. 2009; 71: 117–22. <https://doi.org/10.1016/j.jhin.2008.09.016> PMID: 19013680
29. Rickard CM, Courtney M, Webster J. Central venous catheters: a survey of ICU practices. *J Adv Nurs*. 2004; 48: 247–56. <https://doi.org/10.1111/j.1365-2648.2004.03193.x> PMID: 15488038
30. Alkubati SA, Ahmed NT, Mohamed ON, Fayed AM, Asfour HI. Health care workers' knowledge and practices regarding the prevention of central venous catheter-related infection. *Am J Infect Control*. 2015; 43: 26–30. <https://doi.org/10.1016/j.ajic.2014.09.021> PMID: 25448304
31. Guembe M, Perez-Parra A, Gomez E, Sanchez-Luna M, Bustinza A, Zamora E, et al. Impact on knowledge and practice of an intervention to control catheter infection in the ICU. *Eur J Clin Microbiol Infect Dis*. 2012; 31: 2799–808. <https://doi.org/10.1007/s10096-012-1630-x> PMID: 22565225
32. Sessa A, Di Giuseppe G, Albano L, Angelillo IF. An investigation of nurses' knowledge, attitudes, and practices regarding disinfection procedures in Italy. *BMC Infect Dis*. 2011; 11: 148. <https://doi.org/10.1186/1471-2334-11-148> PMID: 21612613
33. Al-Sayaghi KM. Management of central venous catheters at the intensive care units in Yemen. Survey of practices. *Saudi Med J*. 2011; 32: 275–82. PMID: 21384064
34. Ider BE, Adams J, Morton A, Whitby M, Muugolog T, Lundeg G, et al. Using a checklist to identify barriers to compliance with evidence-based guidelines for central line management: a mixed methods study in Mongolia. *Int J Infect Dis*. 2012; 16: e551–7. <https://doi.org/10.1016/j.ijid.2012.03.006> PMID: 22608032
35. Boersma RS, Schouten HC. Clinical practices concerning central venous catheters in haematological patients. *Eur J Oncol Nurs*. 2010; 14: 200–4. <https://doi.org/10.1016/j.ejon.2009.12.005> PMID: 20117964
36. Snarski E, Mank A, Iacobelli S, Hoek J, Styczynski J, Babic A, et al. Current practices used for the prevention of central venous catheter-associated infection in hematopoietic stem cell transplantation recipients: a survey from the Infectious Diseases Working Party and Nurses' Group of EBMT. *Transpl Infect Dis*. 2015; 17: 558–65. <https://doi.org/10.1111/tid.12399> PMID: 25953418